

# Covering Spaces

in Homotopy Type Theory

Favonia      Robert Harper


Carnegie Mellon University  
{favonia,rwh}@cs.cmu.edu

This material is based upon work supported by the  
National Science Foundation under Grant No. 1116703.



# Homotopy Type Theory (HoTT)

$A$	Type	Space
$a : A$	Term	Point
$f : A \rightarrow B$	Function	Continuous Mapping
$C : A \rightarrow \text{Type}$	Dependent Type	Fibration
$C(a)$		Fiber
$a =_A b$	Identity	Path



# Every type is an $\infty$ -groupoid





# Every type is an $\infty$ -groupoid





# Every type is an $\infty$ -groupoid



# Every type is an $\infty$ -groupoid




# Every type is an $\infty$ -groupoid




$$f : A \longrightarrow B$$

$$a : A \longmapsto b : B$$

$$p : a_1 = a_2 \longmapsto q : b_1 = b_2$$


$$\vdots$$







type

A




type




groupoid

A




type


$A$



groupoid

$\|A\|_1$





# Covering Spaces

Continuously changing families of sets

Classical definition:

A covering space of  $A$  is a space  $C$  together with a continuous surjective map  $p : C \rightarrow A$ , such that for every  $a \in A$ , there exists an open neighborhood  $U$  of  $a$ , such that  $p^{-1}(U)$  is a union of disjoint open sets in  $C$ , each of which is mapped homeomorphically onto  $U$  by  $p$ .

HoTT definition:

$$F : A \longrightarrow \mathbf{Set}$$

Question: Is it correct (up to homotopy)?

# Covering Spaces

$F : A \longrightarrow \text{Set}$

$a : A \longmapsto F(a) : \text{Set}$

$p : a_1 = a_2 \longmapsto \text{iso} : F(a_1) = F(a_2)$

$q : p_1 = p_2 \longmapsto (\text{trivial})$

$\vdots$

# Classification Theorem

Suppose  $A$  is pointed ( $a_0$ ) and connected.

$$F : A \longrightarrow \mathbf{Set}$$

$$\Downarrow$$

$$a_0 : A \longmapsto F(a_0) : \mathbf{Set}$$

$$\text{loop} : a_0 = a_0 \longmapsto \text{auto} : F(a_0) = F(a_0)$$

This is an action of  $\|a_0 = a_0\|_0$  on  $F(a_0)$ .  
 $\|a_0 = a_0\|_0$  is the fundamental group  $\pi_1(A, a_0)$ .



# Classification Theorem

Suppose  $A$  is pointed ( $a_0$ ) and connected.

$$(A \rightarrow \mathbf{Set}) \cong \pi_1(A, a_0)\text{-Set}$$

Pointed ( $a_0$ ) and connected:

$$(a_0 : A) \times ((x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1})$$

Fundamental group  $\pi_1(A, a_0)$ :  $\|a_0 = a_0\|_0$

$$\mathbf{G}\text{-Set}: (X : \mathbf{Set}) \times (\alpha : \mathbf{G} \rightarrow (X \rightarrow X)) \times \\ (\alpha \text{ unit} = \text{id}) \times (\alpha (g_1 \cdot g_2) = \alpha g_1 \circ \alpha g_2)$$

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

$$(A \rightarrow \mathbf{Set}) \simeq \pi_1(A, a_0)\text{-Set}$$

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .


$(A \rightarrow \mathbf{Set}) \simeq \pi_1(A, a_0)\text{-Set}$

$F \mapsto (F(a_0), \star_0, \dots)$

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

$(A \rightarrow \mathbf{Set}) \simeq \pi_1(A, a_0)\text{-Set}$

$F \mapsto (F(a_0), \star_0, \dots)$



transport  $x$  along  $p$  ( $p_* x$ )

$\star : a_1 = a_2 \rightarrow F(a_1) \rightarrow F(a_2)$

$\star_0 : \|a_1 = a_2\|_0 \rightarrow F(a_1) \rightarrow F(a_2)$


( $\star$  for set-truncated paths)

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

$(A \rightarrow \mathbf{Set}) \simeq \pi_1(A, a_0)\text{-Set}$

$F \mapsto (F(a_0), \star_0, \dots)$

$? \longleftarrow (X, \alpha, -, -)$



Idea: formal transports

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

$(A \rightarrow \mathbf{Set}) \simeq \pi_1(A, a_0)\text{-Set}$

$F \mapsto (F(a_0), \star_0, \dots)$

$R_{X,\alpha} \longleftarrow (X, \alpha, -, -)$

$R_{X,\alpha}(a) := X \times \|a_0 = a\|_0$  quotiented by *some relation*  $\sim$ .

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

$(A \rightarrow \mathbf{Set}) \cong \pi_1(A, a_0)\text{-Set}$

$F \longmapsto (F(a_0), \star_0, \dots)$

$R_{X,\alpha} \longleftarrow (X, \alpha, \_ , \_)$

$R_{X,\alpha}(a) := X \times \|a_0 = a\|_0$  quotiented by *some relation*  $\sim$ .

---

**Goal:**  $F = R_{F(a_0), \star_0}$

$F(a) \cong F(a_0) \times \|a_0 = a\|_0$  quotiented by *some relation*  $\sim$ .

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

**Goal:**  $F = R_{F(a_0), \star 0}$

$F(a) \cong F(a_0) \times \|a_0 = a\|_0$  quotiented by *some relation*  $\sim$ .

$$p_{\star 0} x \iff (x, p)$$



Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

**Goal:**  $F = R_{F(a_0), \star 0}$

$F(a) \cong F(a_0) \times \|a_0 = a\|_0$  quotiented by *some relation*  $\sim$ .

$$p_{\star 0} x \iff (x, p)$$

$$x \mapsto (q^{-1}_{\star 0} x, q)?$$

We only have  $\|a_0 = a\|_{-1}$  but need  $q : \|a_0 = a\|_0$ .

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

**Goal:**  $F = R_{F(a_0), \star 0}$

$F(a) \cong F(a_0) \times \|a_0 = a\|_0$  quotiented by *some relation*  $\sim$ .

$$p_{\star 0} x \iff (x, p)$$

$$x \mapsto (q^{-1}_{\star 0} x, q)?$$

We only have  $\|a_0 = a\|_{-1}$  but need  $q : \|a_0 = a\|_0$ .

**Lemma:** If  $(q_1^{-1}_{\star 0} x, q_1) = (q_2^{-1}_{\star 0} x, q_2)$  then  $\|a_0 = a\|_{-1}$  is fine.

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

**Goal:**  $F = R_{F(a_0), \star 0}$

$F(a) \cong F(a_0) \times \|a_0 = a\|_0$  quotiented by *some relation*  $\sim$ .

Wants  $(q_1^{-1} \star_0 x, q_1) = (q_2^{-1} \star_0 x, q_2)$ .

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

**Goal:  $F = R_{F(a_0), \star 0}$**

$F(a) \cong F(a_0) \times \|a_0 = a\|_0$  quotiented by *some relation*  $\sim$ .

Wants  $(q_1^{-1} \star_0 x, q_1) = (q_2^{-1} \star_0 x, q_2)$ .

$(\alpha \text{ loop } x, p) \sim (x, \text{loop } \blacksquare p)$

Intuition:  $p \star_0 (\text{loop } \star_0 x) = (\text{loop } \blacksquare p) \star_0 x$

$$\begin{aligned} (q_1^{-1} \star_0 x, q_1) &= (q_1^{-1} \star_0 x, (q_1 \blacksquare q_2^{-1}) \blacksquare q_2) \\ &= ((q_1 \blacksquare q_2^{-1}) \star_0 (q_1 \star_0 x), q_2) = (q_2^{-1} \star_0 x, q_2) \end{aligned}$$

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

$(A \rightarrow \mathbf{Set}) \simeq \pi_1(A, a_0)\text{-Set}$

$F \mapsto (F(a_0), \star_0, \dots)$

$R_{X,\alpha} \longleftarrow (X, \alpha, -, -)$

$R_{X,\alpha}(a) := X \times \|a_0 = a\|_0$  quotiented by  
 $(\alpha \text{ loop } x, \text{ path}) \sim (x, \text{ loop } \blacksquare \text{ path})$

Suppose  $a_0 : A$  and  $(x : A) \rightarrow (y : A) \rightarrow \|x = y\|_{-1}$ .

$(A \rightarrow \mathbf{Set}) \simeq \pi_1(A, a_0)\text{-Set}$

$F \mapsto (F(a_0), \star_0, \dots)$

$R_{X,\alpha} \longleftarrow (X, \alpha, \_ , \_)$

$R_{X,\alpha}(a) := X \times \|a_0 = a\|_0$  quotiented by  
 $(\alpha \text{ loop } x, \text{ path}) \sim (x, \text{ loop } \blacksquare \text{ path})$

---

The other round trip is easy.

$(G\text{-sets} \rightarrow \text{covering spaces} \rightarrow G\text{-sets})$

# Summary

- A simple formulation:  $A \rightarrow \text{Set}$ .
- Type equivalence of  $A \rightarrow \text{Set}$  and  $\pi_1(A)\text{-Set}$ .

# Summary

- A simple formulation:  $A \rightarrow \text{Set}$ .
- Type equivalence of  $A \rightarrow \text{Set}$  and  $\pi_1(A)\text{-Set}$ .

# Notes

- Other theorems (universal coverings, categories).
- Fibers need not to be decidable types.
  - ☞ “path-constant” spaces, not just discrete ones?
- $A \rightarrow \text{Groupoid}$ ?



# Summary

- A simple formulation:  $A \rightarrow \text{Set}$ .
- Type equivalence of  $A \rightarrow \text{Set}$  and  $\pi_1(A)\text{-Set}$ .

# Notes

- Other theorems (universal coverings, categories).
- Fibers need not to be decidable types.
  - ☞ “path-constant” spaces, not just discrete ones?
- $A \rightarrow \text{Groupoid}$ ?

# Thank you

**Acknowledgements:** Carlo Angiuli, Steve Awodey, Andrej Bauer, Spencer Breiner, Guillaume Brunerie, Daniel Grayson, Chris Kapulkin, Nicolai Kraus, Peter LeFanu Lumsdaine and Ed Morehouse